

## DISCHARGE INSTRUCTIONS FOLLOWING UFE

---

### Activities:

What a relief to finally have the UFE procedure behind you! For the first several days following the procedure many women describe feeling like they have the flu ... achy and tired. For the first week, you are encouraged to take it easy with no heavy lifting, no vigorous exercise, and no sexual intercourse. If you typically use tampons, please use another means of feminine protection for the first month following UFE. You should feel a little better each day and will likely return to your usual activities in 7-10 days, if not sooner. Remember, any pain/discomfort you may experience represents the process of fibroid degeneration (which is good as it means the fibroids are dying) and pales in comparison to the pain after myomectomy or hysterectomy. If you overdo it your body will tell you and it may set you back for a day or two.

### What to Watch For:

Call your Interventional Radiologist (not your gynecologist) at 889-5749 with any problems or questions. If you need an immediate response but get our voicemail, hang up and call our answering service at **455-3503** and ask them to page your Interventional Radiologist. In particular, **CALL IF** you have a **temperature greater than 100.5°** or **pain that is not being relieved** by the medication. It is not unusual to develop a vaginal discharge, sometimes even a mini-period, and you may pass small fragments of tissue. Call if you develop a **green or yellow vaginal discharge**. The groin puncture site may bruise and be sore for a few days, but it should not be extremely painful or develop a **large knot**. Call if you develop **calf pain, swelling, or shortness of breath**.

### Medications:

**Motrin**, 800-*mg*, three times per day for seven days ... take it for the full seven days, even if you feel completely better. If the Motrin upsets your stomach, take it with food or Maalox. It does not matter what form of ibuprofen (the generic name for Motrin) you take, as long as you take 800-*mg* three times daily. The over-the-counter ibuprofen (Advil and others) is usually 200-*mg* per tablet, so you could take four at a time.

If you have pain that the Motrin does not relieve, add **Vicodin** (a mild narcotic). You may take as many as two pills every four hours, but you do not have to take it at all if you so wish. You are taking too much if it makes you sleepy. The Vicodin is a supplement to the Motrin, **NOT** a substitute for it. Don't hesitate to take both, if needed. Do not drink alcohol or drive while taking the Vicodin.

The Vicodin can make you constipated, which you really don't want to happen. When you fill the other prescriptions, purchase a box of **Surfak**, an over-the-counter stool softener. Take one dose three times a day until your bowel function has returned to normal. Do **NOT** take any iron supplements for the first ten days after UFE as they aggravate the constipation.

### Follow-Up:

If not already arranged following your UFE, please call 889-5749 to schedule a follow-up appointment with your Interventional Radiologist at the DePaul Medical Building / VBGH / MIH office. Your appointment is scheduled for \_\_\_\_\_ @ \_\_\_\_\_.

Call our office in three months following the UFE to schedule a brief follow-up appointment. Contact our office to schedule a follow-up MRI of the pelvis in six months following the UFE, to see how your uterus and fibroid(s) have responded to the embolization. Please schedule this appointment at the same hospital where you had your pre-UFE MRI.

Congratulations! .... you are well on your way to life after fibroids!